

CONTRIBUTIONS TO THE DOLICHOPODIDAE (DIPTERA) FAUNA OF THE MARMARA REGION OF TURKEY WITH NEW RECORDS

ALPER TONGUÇ

Molecular Biology and Genetics Department, Faculty of Science, Muğla Sıtkı Koçman University,
48000, Muğla, Turkey; Tel: +90 252 2115569
E-mail: alpertonguc@mu.edu.tr

Abstract

Field surveys were carried out in the provinces of Balıkesir, Bilecik, Bursa, Çanakkale, Edirne, İstanbul, İzmit, Kırklareli, Sakarya, Tekirdağ and Yalova in the Marmara region of Turkey between 2009 and 2010. As a result of identification of the collected specimens, 17 genera and 31 species, belonging to Diaphorinae, Dolichopodinae, Hydrophorinae, Rhaphininae, Sciapodinae, Sympycninae, Xanthochlorinae subfamilies, were determined. Two species (*Hercostomus plagiatus* Loew, 1857 and *Poecilobothrus nobilitatus* Linnaeus, 1767) are new records for Turkey dolichopodid fauna.

KEY WORDS: Dolichopodidae, fauna, Marmara region, Turkey, new records

Introduction

Dolichopodidae (long-legged flies) is one of the biggest families in the Diptera order. The body size ranges between 1-9 mm. They are known for the metallic-like appearance of the body, arista-type antenna, wing venation and male genitalia. Almost all long legged-flies are polyphagous predators feeding on various fine invertebrates including some pests. The adult flies live in different habitat types such as aquatic and semiaquatic areas, marshlands, woodlands, coastal dunes, grasslands and heathland (Grichanov, 2007).

This family contains 257 extant and 30 fossil genera, and 7 945 valid species. About 1 553 species are known from the Palearctic region (Grichanov, 2017). A checklist of Dolichopodidae in Turkey that contains one subspecies and 187 species was published in 2016 (Tonguc et al., 2016a). Fourteen new species were added for the country after 2016 and three of them were recorded for the European part of Turkey (Tonguç et al. 2016b; Küçükberber et al. 2017; Grichanov & Ahmadi, 2017; Naglis & Negrobov 2017). So far, a total of 37 species have been known from the entire Marmara region, 30 species from the European part

(Kechev *et al.*, 2020) and 12 species from other parts of the region (Grichanov *et al.* 2007; Tonguc *et al.*, 2009; Grichanov and Tonguc, 2010a; 2010b; Tonguc *et al.*, 2010). The Dolichopodidae fauna of Turkey includes 203 species. This study aims to contribute to records of the Dolichopodidae fauna of the Marmara region and Turkey as a whole.

Materials and Methods

This study was carried out in four sub-regions of the Marmara region (Çatalca-Kocaeli, Ergene, Yıldız Mountains and South Marmara) between July 2008 and December 2010. The research area includes İstanbul, Edirne, Kırklareli, Tekirdağ, Yalova, İzmit, Sakarya, Bilecik, Bursa, Balıkesir and Çanakkale provinces. Specimens were collected with an entomological hand net from different habitat types and preserved in insect envelops or 70% ethyl alcohol in the research field. Some of the Dolichopodidae specimens brought to the laboratory were pinned, and others were preserved in different sizes of glasses or plastic jars. All of the specimens are stored in insect collection boxes in the Faculty of Science, Department of Biology, Laboratory of Zoology of Muğla Sıtkı Koçman University. A faunistic list of species distribution belonging to the provinces, Palearctic and Turkey distributions are given in the Results. Palearctic distribution is presented by Yang *et al.* (2006) and Grichanov (2007) according to following country abbreviations: Afghanistan (AF), Albania (AL), Algeria (DZ), Armenia (AM), Azerbaijan (AZ), Austria (AT), Belarus (BY), Belgium (BE), Bosnia-Herzegovina (BA), Bulgaria (BG), Croatia (HR), Czech Republic (CZ), Denmark (DK), Egypt (EG), England (EN), Estonia (EE), Finland (FI), France (FR), Georgia (GE), Greece (GR), Germany (DE), Holland (NL), Hungary (HU), Iraq (IQ), Iran (IR), Ireland (IE), Israel (IL), Italy (IT), Kazakhstan (KZ), Kyrgyzstan (KG), Lithuania (LT), Luxemburg (LU), Macedonia (MK), Moldova (MD), Morocco (MA), Norway (NO), Poland (PL), Portugal (PT), Romania (RO), Russia (RU), Slovakia (SK), Slovenia (SI), Spain (ES), Sweden (SE), Switzerland (CH), Syria (SY), Tajikistan (TJ), Tunisia (TN), Turkmenistan (TM), Turkey (TR), Ukraine (UA), Uzbekistan (UZ).

Results

Faunistic List

Diaphorinae Schiner, 1864

***Argyra argyria* (Meigen, 1824)**

Material examined: Kırklareli, Pınarhisar, İslambeyli, 41°41' N, 27°37' E, 342 m, 28.07.2009, 1 ♂.

Palearctic Distribution: TR, AT, BE, BY, CH, CZ, DE, DK, EN, ES, FI, FR, GR, HR, HU, IT, SE, MA, MD, NL, NO, PL, RO, RU, SK, UA.

Turkey Distribution: Afyonkarahisar, Kütahya, Uşak.

***Diaphorus vitripennis* Loew, 1859**

Material examined: Bursa, Osmangazi, Demirtaş Village, Baraj Lake, 40°17' N, 29°06' E, 135 m, 27.04.2009, 1 ♂.

Palearctic Distribution: TR, AT, CH, DZ, FR, HU, IT, KZ, PT, RO, UZ.

Turkey Distribution: Aydın (Buharkent), Kars.

Dolichopodinae Latreille, 1809

***Dolichopus griseipennis* Stannius, 1831**

Material examined: Bursa, Osmangazi, Uludağ Road 5 km, 40°08' N, 29°01' E, 875 m, 19.05.2009, 1 ♂; Uludağ, Hüseyinalan Village, 40°07' N, 29°01' E, 900 m, 06.08.2010, 1 ♂; Uludağ 2. Oteller Place, 40°06' N, 29°09' E, 1780 m, 06.08.2010, 1 ♀; Uludağ 2. Oteller Place, 40°06' N, 29°09' E, 1780 m, 07.08.2010, 1 ♂, 2 ♀♀; Uludağ Road, 40°09' N, 29°01' E, 755 m, 06.08.2010, 2 ♂♂, 2 ♀♀; Uludağ M. P. Road, 40°08' N, 29°01' E, 875 m, 06.08.2010, 1 ♂; İnegöl, Güneykestane, 39°56' N, 29°43' E, 550 m 03.08.2010, 1 ♂; Karacabey, Yenikaraağaç village, 40°13' N, 28°37' E, 45 m, 07.08.2010, 2 ♂♂, 3 ♀♀; Mustafakemalpaşa, Muradiyesarnıcı Village, 39°55' N, 28° 23' E, 247 m, 09.07.2008, 3 ♂♂, Çanakkale, Biga, Çelikgürü Village, 40°19' N, 27°02' E, 46 m, 25.04.2010, 1 ♀; Bayramiç, Çandır, 39°46' N, 26°34' E, 146 m, 26.07.2009, 1 ♂; Yenice, 39°56' N, 27°13' E, 281 m, 26.07.2009, 3 ♂♂, 1 ♀; Kocaeli, Gebze, Yağcılar Village, 19.05.2010, 1 ♂, 3 ♀♀; Sakarya, Adapazarı, Çaybaşıyeniköy, 40°40' N, 30°27' E, 75 m, 17.05.2010, 1 ♀.

Palearctic Distribution: TR, AM, AT, AZ, BA, BE, BG, CH, CZ, DE, DK, DZ, EE, EN, ES, FI, FR, GE, GR, HR, HU, IE, KZ, LU, MA, MK, NL, NO, PL, RO, RU, SE, SI, SK, TN.

Turkey Distribution: Burdur, Muğla, Sinop.

***Dolichopus salictorum* Loew, 1871**

Material examined: Bursa, Osmangazi, Uludağ 2. Oteller Place, 40°06' N, 29°09' E, 1780 m, 07.08.2010, 4 ♂♂, 1 ♀, İnegöl, 39°52' N, 29°39' E, 1146 m, 03.08.2010, 1 ♂, 1 ♀.

Palearctic Distribution: TR, BG, CZ, HU, IT, PL, RO, SK.

Turkey Distribution: Adiyaman (Naglis, 2011).

***Dolichopus signifer* Haliday, 1832**

Material examined: Çanakkale, Eceabat, Küçükanaftalar, 40°16' N 26°17' E, 25 m, 11.04.2010, 1 ♂; Balıkesir, Edremit, Altınoluk, Kaz Mountains, Güvercin Place, 39°41' N 26°47' E, 1147 m, 27.08.2009, 1 ♀.

Palearctic Distribution: TR, AM, AT, BE, BG, CH, CZ, DE, EN, ES, FR, GE, GR, HU, IE, IT, KZ, MA, NL, NO, PL, RO, RU, SE, SK, TJ, TM, UA, UZ.

Turkey Distribution: Burdur, Edirne.

***Hercostomus fulvicaudis* (Walker, 1851)**

Material examined: Bursa, İnegöl, Güneykestane, 39°56' N 29°43' E, 550 m 03.08.2010, 7 ♀♀.

Palearctic Distribution: TR, AT, BE, CZ, EN, FR, DE, NL, PL, RO, SK, SE, TM, TJ, UA.

Turkey Distribution: Afyonkarahisar, Kütahya.

***Hercostomus gracilis* (Stannius, 1831)**

Material examined: Balıkesir, Dursunbey, 39°36' N 28°38' E, 595 m, 15.07.2010, 1 ♂, 1 ♀; Edremit, Kaz Mountains, 39°41' N 27°09' E, 560 m 26.08.2009, 1 ♂; Çınarlıhan Place, 39°41' N 27°10' E, 660 m, 08.08.2010, 7 ♂♂; Mehmetalan Village, 39°40' N 26°58' E, 335 m, 12.07.2008, 3 ♂♂, 2 ♀♀; Bursa,

Osmangazi, Karaisalı Village, 40°02' N 29°05' E, 815 m, 25.08.2009, 1 ♂; Uludağ- Soğukpınar Village, 40°03' N 29°07' E, 1025 m 25.07.2009, 1 ♂; Uludağ, Soğukpınar-Keles Road 15.Km, 39°58' N 29°13' E, 965 m 25.07.2009, 1 ♂, 1 ♀.

Palearctic Distribution: TR, AM, AT, BE, BG, CZ, DK, EN, ES, FR, DE, GR, HU, IT, NL, PL RU, SE, TJ, TM, UA.

Turkey Distribution: Aydın, Burdur (Grichanov et al., 2007).

***Hercostomus longiventris* (Loew, 1857)**

Material examined: Sakarya, Geyve, Doğançay village, 40°36' N 30°19' E, 45 m, 25.10.2008, 1 ♂.

Palearctic Distribution: TR, AT, BA, BE, CH, CZ, DE, FR, GR, HR, IT, NL, HU, MA, MK, PL, RO, RU, SI, TJ.

Turkey Distribution: Muğla, Artvin (Grichanov vd., 2007).

***Hercostomus plagiatus* (Loew, 1857)**

Material examined: Bursa, Osmangazi, Uludağ National Park, Sarıalan, 40°08' N 29°06' E, 1620 m, 06.08.2010, 1 ♂.

Palearctic Distribution: TR, AT, BE, CH, CZ, DE, EN, ES, FR, HU, IT, MK, NL, RO, SI, TN.

Turkey Distribution: New records for Turkey.

***Hercostomus phoebus* Parent, 1927**

Material examined: Bursa, Osmangazi, Uludağ, Soğukpınar-Keles Road 15. Km, 39°58' N 29°13'E, 965 m 25.07.2009, 1 ♂, 1 ♀; Kestel, Hüseyinalan village, 39°45' N 29°12' E, 746 m, 25.07.2009, 2 ♂♂, 2 ♀♀.

Palearctic Distribution: TR, AM.

Turkey Distribution: Ankara, Antalya, Burdur, Muğla (Parent, 1927; Hollis, 1963; Grichanov vd., 2007).

***Hercostomus rusticus* (Meigen, 1824)**

Material examined: Çanakkale, Bayramiç, Evciler town, Kaz Mountains, 39°41' N 26°47' E, 1265 m, 19.08.2008, 1 ♂.

Palearctic Distribution: TR, AM, AT, BA, BE, BG, BY, CH, CZ, DE, EE, ES, FR, GE, GR, HR, IT, KZ, MK, MN, NL, PL, RO, RU, SI, SK, UK.

Turkey Distribution: Rize (Naglis, 2011).

***Hercostomus thraciensis* Kechev & Negrobov, 2015**

Material examined: Bursa, Mustafakemalpaşa, Yalıntaş village, 39°58' N 28°22' E, 253 m, 28.04.2009, 1 ♂.

Palearctic Distribution: TR, BG.

Turkey Distribution: Manisa

***Ortochile nigrocoerulea* Latreille, 1809**

Material examined: Bursa, Mustafakemalpaşa, Yalıntaş Village, 39°58' N 28°22' E, 253 m, 28.04.2009, 3 ♂♂, 4 ♀♀; Balıkesir, Gönen, Gündoğan Village, 40°09' N 27°38' E, 45 m, 25.04.2010, 1 ♂, 3 ♀♀; Bilecik,

Osmaneli, Gaziler Village, 40°24' N 29°56' E, 310 m, 16.05.2010, 1 ♂, 2 ♀♀; Çanakkale, Ayvacık, Kocaköy, 39°29' N 26°09' E, 265 m, 10.04.2010, 1 ♂, 1 ♀.

Palearctic Distribution: TR, AT, BE, SE, DK, DZ, EN, ES, FR, GR, HR, HU, IL, IT, MK, PL, TN.

Turkey Distribution: Aydın, Kahramanmaraş.

***Poecilobothrus nobilitatus* (Linneaus, 1767)**

Material examined: Bursa, Osmangazi, Uludağ- Soğuksınar Village, 40°03' N 29°07' E, 1025 m 25.07.2009, 2 ♂♂; İnegöl, Eskikaracaköy, Domaniç Road, 39°54' N 29°40' E, 1125 m, 24.07.2009, 1 ♂.

Palearctic Distribution: TR, AT, BE, CH, CZ, DE, DK, EN, ES, FR, HU, IE, IT, LU, MK, NL, PL, RO, RU, SE, SK, UK.

Turkey Distribution: New records for Turkey.

***Poecilobothrus principalis* (Loew, 1861)**

Material examined: Bilecik, Bozüyüük, Muratdere, 39°53' N 29°53' E, 875 m, 05.08.2010, 1 ♂.

Palearctic Distribution: TR, AT, BE, BG, CZ, DE, EN, ES, FR, HU, IL, IT, NL, PL, RO, RU, UA.

Turkey Distribution: Kütahya.

***Poecilobothrus regalis* (Meigen, 1824)**

Material examined: Balıkesir, Manyas, Haydar Village, 40°04' N 28°00' E, 29 m, 03.06.2009, 4 ♂♂, 2 ♀♀; İvrindi, Soğanbüyük Village, Güngörmez Stream, 39°36' N 27°32' E, 207 m, 23.6.2009, 5 ♂♂, 1 ♀; Bilecik, Bozüyüük, 39°51' N 29°58' E, 850 m, 07.07.2010, 3 ♂♂, 2 ♀♀; Gölpazarı, Karaahmetler Village, 40°19' N 30°27' E, 740 m, 17.07.2008, 2 ♂♂, 3 ♀♀; Muratdere, 39°53' N 29°53' E, 875 m, 05.08.2010, 12 ♂♂, 2 ♀♀; Bursa, Mustafakemalpaşa, Sinansarnıcı Village, 40°03' N 28°40' E, 466 m, 24.06.2009, 4 ♂♂, 3 ♀♀; Nilüfer, Kayapa Village, 40°10' N 28°51' E, 128 m, 24.06.2009, 17 ♂♂, 5 ♀♀; İznik, Derbent Village, 40°21' N 29°40' E, 275 m, 18.07.2008; 5 ♂♂, 13 ♀♀; 26.06.2009, 7 ♂♂; Osmangazi, Uludağ, Soğuksınar-Keles Road 15.Km, 39°58' N 29°13' E, 965 m 25.07.2009, 1 ♂, 1 ♀; Çanakkale, Bayramiç, Ağaçköy, 39°47' N 26°34' E, 123 m, 26.07.2009, 2 ♀♀; Kocaeli, Karamürsel, Valide Köprü, 40°34' N 29°31' E, 130 m, 05.06.2009, 5 ♂♂, 2 ♀♀; Sakarya, Taraklı, Çamtepe Village, 40°21' N 30°31' E, 460 m, 17.07.2008, 3 ♂♂; Geyve, Çamlık Village, 40°34' N 30°21' E, 436 m, 06.06.2009, 5 ♂♂; İstanbul, Silivri, Değirmenköy, 41°07' N 28°01' E, 110 m, 16.07.2008, 4 ♂♂, 6 ♀♀; Yalova, Altınova, Fevziye Village, 40°38' N 29°30' E, 45 m, 17.07.2008, 3 ♀♀.

Palearctic Distribution: TR, AT, BE, BG, CZ, DE, FR, GE, GR, IR, IT, MK, RO, RU, SK, UA, UZ.

Turkey Distribution: Burdur, Denizli, Isparta, Muğla (Grichanov vd., 2007), Antalya (Parvu ve Popescu-Mirceni, 2006; Grichanov vd., 2007).

***Sybistroma sphenopterus* (Loew, 1859)**

Material examined: Bursa, Osmangazi, Uludağ Road 7. Km, 40°09' N 29°01' E, 780 m, 09.07.2008, 1 ♂; Kocaeli, İzmit, Maşukkiye Village, Şelale Place, 40°41' N 30°07' E, 250 m, 23.08.2009, 2 ♂♂.

Palearctic Distribution: TR, AT, CZ, DEHU, IT, PL, RO.

Turkey Distribution: Çanakkale (Tonguç vd., 2009), Kars (Naglis, 2011).

Hydrophorinae Lioy, 1864

***Hydrophorus balticus* (Meigen, 1824)**

Material examined: Balıkesir, Edremit, Kaz Mountains, 39°41' N 27°09' E, 560 m 26.08.2009, 3 ♀♀; Hanlar Village, 39°41' N 27°09' E, 625 m, 11.07.2008, 2 ♂♂; Bursa, Osmangazi, Uludağ National Park, 1.Oteller Place, 40°06' N 29°08' E, 1790 m, 25.06.2009, 7 ♂♂, 18 ♀♀; Çanakkale, Lapseki, Kocababaşlar Village, 40°17' N 26°50' E, 565 m, 26.04.2010, 2 ♂♂, 1 ♀; Bayramiç, Evciler Town, Kaz Mountains, Düden Place, 39°41' N 26°48' E, 1265 m, 27.08.2009, 9 ♂♂, 8 ♀♀; Kocaeli, İzmit, Maşukkiye Village, Kartepe Road 13.Km, 40°39' N 30°07' E, 1184 m, 23.08.2009, 1 ♂, 1 ♀; Kartepe, 40°39' N 30°07' E, 1190 m, 09.07.2010, 5 ♂♂, 6 ♀♀.

Palearctic Distribution: TR, AF, AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, EN, ES, FI, FR, GE, GR, HU, IE, IL, IT, MA, NL, NO, PL, RO, RU, SE, UA.

Turkey Distribution: Aydin, Denizli, Isparta, Rize, Antalya, Manisa.

***Hydrophorus praecox* (Lehmann, 1822)**

Material examined: Bilecik, Bozüyük, 39°51' N 29°58' E, 850 m, 07.07.2010, 1 ♀.

Palearctic Distribution: TR, AT, BE, BG, CH, CN, CY, CZ, DE, DK, EE, EG, EN, ES, FI, FR, GE, HU, IE, IL, IQ, IR, KZ, NL, NO, PL, RO, RU, SE, SK, UA.

Turkey Distribution: Bolu, Burdur, Antalya.

***Liancalus virens* (Scopoli, 1763)**

Material examined: Balıkesir, Edremit, Kaz mountains, 39°41' N 27°09' E, 650 m, 08.08.2010, 1 ♂, 1 ♀; Bilecik, Bozüyük, Kozpinar Village, 39°54' N 29°47' E, 540m, 21.08.2009, 1 ♀; Bursa, Osmangazi, Uludağ National Park, Sarialan, 40°08' N 29°06' E, 1620 m, 06.08.2010, 1 ♀; Çanakkale, Gökçeada, Aydıncık, 40°10' N 25°54' E, 170 m, 13.07.2010, 1 ♂, 1 ♀; Kocaeli, Karamürsel, Oluklu Village, 40°39' N 29°35' E, 375 m, 04.05.2009, 3 ♂♂, 1 ♀.

Palearctic Distribution: TR, AT, BE, BG, CH, CY, CZ, DE, DK, EN, ES, FI, FR, GE, GR, HU, IE, IL, KG, KZ, LU, SE, MA, NL, NO, PL, RO, RU, SK, TJ, TN, UA.

Turkey Distribution: Muğla, Manisa.

***Orthoceratium lacustre* (Scopoli, 1763)**

Material examined: Çanakkale, Bayramiç, Evciler Town, Kaz Mountains, 6. Km, 39°43' N 26°47' E, 740 m, 19.08.2008, 1 ♀; Kaz Mountains, 16. Km, 39°42' N 26°48' E, 1185 m, 13.07.2008, 3 ♂♂, 6 ♀♀.

Palearctic Distribution: TR, AL, AT, AZ, BE, BG, CY, DE, DK, EN, ES, IE, FI, FR, GR, IL, IT, NL, PT, TN, UA.

Turkey Distribution: Afyonkarahisar, Kütahya.

***Scellus notatus* (Fabricius, 1781)**

Material examined: Balıkesir, Gönen, Gündoğan Village, 40°09' N 27°38' E, 45 m, 25.04.2010, 1 ♂; Bursa, Osmangazi, Uludağ National Pak 10. Km, 40°06' N 29°06' E, 1740 m, 25.07.2009, 1 ♀; Bakacak Place, 40°07' N 29°08' E, 1745 m, 25.06.2009, 2 ♀♀; Çanakkale, Yenice, Umurlar Village, 39°53' N 27°22' E, 220 m, 26.04.2010, 1 ♂.

Palearctic Distribution: TR, AT, BE, BG, DE, DK, EN, FR, GR, HU, IE, IT, NL, PT, RO, RU, SE, SK, UA.

Turkey Distribution: Afyonkarahisar, Manisa.

Rhaphininae Bigot, 1852

***Rhaphium appendiculatum* Zetterstedt, 1849**

Material examined: Bursa, Osmangazi, Uludağ- Soğuksınar Village, 40°03' N 29°07' E, 1025 m 25.07.2009, 1 ♂, 1 ♀; Çanakkale, Bayramiç, Evciler Town, Kaz Mountains, 11. Km, 39°42' N 26°47' E, 880 m, 27.08.2009, 1 ♂.

Palearctic Distribution: TR, AF, AL, AT, BG, CZ, DE, DK, EN, ES, FI, FR, GE, GR, HU, IE, IR, IT, MA, NL, PL, RO, RU, SE, UA.

Turkey Distribution: Antalya, İzmir, Manisa.

***Rhaphium calliginosum* Meigen, 1824**

Material examined: Bilecik, Söğüt, Dereboyu Village, 39°59' N 30°08' E, 1005 m, 02.05.2009, 1 ♂.

Palearctic Distribution: TR, AL, AM, AT, BE, BG, CH, CZ, DE, DK, EE, EN, ES, FI, FR, GR, HU, IE, IL, IT, MA, NL, NO, SK, PL, RO, RU, SE, SY, UA.

Turkey Distribution: Burdur, İzmir.

Sciapodinae Becker, 1917

***Sciapus bellus* (Loew, 1873)**

Material examined: Sakarya, Karasu, Limandere Village, 40°59' N 30°36' E, 18 m, 03.05.2009, 1 ♂.

Palearctic Distribution: TR, AT, CH, CZ, DE, GR, HU, IT, PL, RO, SK.

Turkey Distribution: Sakarya.

Sympycninae Aldrich, 1905

***Campsicnemus curvipes* (Fallén, 1823)**

Material examined: Bursa, Osmangazi, Uludağ- Soğuksınar Village, 40°03' N 29°07' E, 1025 m 25.07.2009, 1 ♂, 2 ♀♀; Sakarya, Kaynarca, Kulaklı Village, 41°04' N 30°23' E, 17 m, 03.05.2009, 1 ♂; Taraklı, Çamtepe Village, 40°21' N 30°31' E, 460 m, 17.07.2008, 1 ♂.

Palearctic Distribution: TR, AL, AT, AM, AZ, BE, BG, BY, CH, CZ, DE, DK, EE, EN, ES, FI, FR, GR, HU, IE, IT, LU, MA, MK, NL, NO, PL, RO, RU, SE, SI, SK, UA.

Turkey Distribution: Bolu, Antalya, Manisa.

***Campsicnemus magius* (Loew, 1845)**

Material examined: Bursa, Mustafakemalpaşa, Muradiyesarnıç Village, 39°55' N 28°23' E, 265 m, 20.10.2008, 1 ♂.

Palearctic Distribution: TR, AT, BE, BG, CZ, DE, EN, ES, FR, HU, IL, IT, NL, RO, RU, SI, TJ, TM, UA.

Turkey Distribution: Sakarya (Korucuk).

***Sympycnus pulicarius* (Fallén, 1823)**

Material examined: Bursa, Osmangazi, Uludağ Road 5. Km, 40°08' N 29°01' E, 875 m, 19.05.2009, 1 ♂; Uludağ- Soğuksınar Village, 40°03' N 29°07' E, 1025 m 25.07.2009, 1 ♂; Çanakkale, Yenice, Kalkım Town, Kaz Mountains, 39°41' N 27°09' E, 625 m, 17.10.2008, 3 ♂♂, 3 ♀♀.

Palearctic Distribution: TR, AT, BE, BG, CH, CZ, DE, DK, EE, EN, ES, FI, FR, GR, HU, IE, IT, LU, NO, NL, PL, RU, RO, SE, SK, UA.

Turkey Distribution: Aydın, Muğla, Van.

***Syntormon denticulatus* (Zetterstedt, 1843)**

Material examined: Çanakkale, Ayvacık, Gülpınar Tuzla Village, 39°34' N 26°10' E, 12 m, 10.04.2010, 1 ♀; Eceabat, Küçükanaftalar, 40°16' N 26°17' E, 25 m, 11.04.2010, 1 ♀; Bursa, Mustafakemalpaşa, Muradiyesarnıç Village, 39°55' N 28°23' E, 265 m, 20.10.2008, 1 ♂.

Palearctic Distribution: TR, AF, AM, BG, DE, EE, FR, IL, IT, Middle Asia, PL, RO, RU, UA.

Turkey Distribution: Muğla, Manisa.

***Syntormon pallipes* (Fabricius, 1794)**

Material examined: Balıkesir, Edremit, Altınluk, Kaz Mountains, Güvercin Place, 39°41' N 26°47' E, 1147 m, 27.08.2009, 16 ♂♂, 9 ♀♀; Bilecik, Bozüyüklü, Sarıdayı Village, 39°55' N 29°48' E, 908 m, 26.06.2009, 1 ♂, 1 ♀; Bursa, Osmangazi, Uludağ National Park, Bakacak stream, 40°07' N 29°08' E, 1745 m, 24.08.2009, 5 ♂♂, 5 ♀♀; Çanakkale, Bayramiç, Evciler Town, Kaz Mountains, 16. Km, 39°42' N 26°48' E, 1185 m, 13.07.2008, 10 ♂♂, 9 ♀♀; Kocaeli, Karamürsel, Valide Köprü, 40°34' N 29°31' E, 130 m, 05.06.2009, 1 ♀; Sakarya, Taraklı, Çamtepe Village, 40°21' N 30°31' E, 460 m, 17.07.2008, 7 ♂♂.

Palearctic Distribution: TR, AL, AF, AT, BE, BG, CH, CN, CZ, DE, DK, EE, EG, EN, ES, FI, FR, GE, GR, HU, IE, IL, IQ, IR, IS, IT, MA, NL, NO, PL, PT, RO, RU, SE, SI, SK, TJ, UA, UZ.

Turkey Distribution: Sakarya (Korucuk), Isparta, Burdur, Denizli, Muğla, Antalya, Bergama, Adiyaman, Ankara, Van, Hakkari, Manisa.

Xanthochlorinæ Loew, 1857

***Xanthochlorus tenellus* (Wiedemann, 1817)**

Material examined: Çanakkale, Bayramiç, Evciler Town, Kaz Mountains, Düden Place, 39°41' N 26°47' E, 1265 m, 19.08.2008, 1 ♂; Sakarya, Akyazı, Sülüklü Lake Natural Reserve, 40°31' N 30°52' E, 1062 m, 22.08.2009, 1 ♀.

Palearctic Distribution: TR, AT, BE, BG, CH, CZ, DE, DK, EE, EN, FI, FR, GE, HU, IE, IT, LT, NL, NO, PL, RO, RU, SE, SK, UA.

Turkey Distribution: Kütahya.

Discussion

Seven subfamilies, 17 genera and 31 species were determined as a result of identification of collected specimens from the research area. *Hercostomus plagiatus* and *Poecilobothrus nobilitatus* are new records for Turkey, whereas 14 species are new for the Marmara region. Fifty-four species are now known from the Marmara region with these additional records (Table I). In Table 1, new records for the investigated provinces are marked with an asterisk (*) and new records for Turkey are marked with a double asterisk (**) in faunistic remarks.

Respectively 8, 7, 20, 13, 1, 6, 8 and 1 species from Balıkesir, Bilecik, Bursa, Çanakkale, İstanbul, Kocaeli, Sakarya and Yalova were collected. Thus, 8 species are known from Balıkesir, 7 from Bilecik, 24 from Bursa, 20 from Çanakkale, 4 from Edirne, 1 from İstanbul, 23 from Kırklareli, 7 from Kocaeli, 11 from Sakarya, 7 from Tekirdağ and 1 from Yalova (Fig 1).

Table I. Distributions of identified species by provinces in Marmara Region.

Species	Balıkesir	Bilecik	Bursa	Çanakkale	Edirne	İstanbul	Kırklareli	Kocaeli	Sakarya	Tekirdağ	Yalova
<i>Argyra argentina</i> (Meigen, 1824)						X					
<i>A. argyria</i> (Meigen, 1824)									X*		
<i>A. diaphana</i> (Fabricius, 1775)					X						
<i>Chrysotus pulchellus</i> Kowarz, 1874									X		
<i>Diaphorus pilosibius</i> Negrobov et Maslova, 2005									X		
<i>D. vitripennis</i> Loew, 1859			X*		X*						
<i>Dolichopus griseipennis</i> Stannius, 1831		X*		X*			X	X*	X*		
<i>D. latilimbatus</i> Macquart, 1827									X		
<i>D. nubilus</i> Meigen, 1824								X			
<i>D. plumipes</i> (Scopoli, 1763)								X			
<i>D. salictorum</i> Loew, 1871			X*								
<i>D. signifer</i> Haliday, 1832	X*			X*			X				
<i>Gymnopternus celer</i> (Meigen, 1824)					X						
<i>Hercostomus fulvicaudis</i> (Walker, 1851)				X*							
<i>H. gracilis</i> (Stannius, 1831)	X*		X*				X				
<i>H. libanicola</i> Parent, 1938					X		X				
<i>H. longiventris</i> (Loew, 1857)									X*		
<i>H. plagiatus</i> (Loew, 1857)			X*								
<i>H. phoebus</i> Parent, 1927			X*								
<i>H. rusticus</i> (Meigen, 1824)					X*						
<i>H. strobianus</i> Becker, 1917							X				
<i>H. thraciensis</i> Kechev & Negrobov, 2015	X*		X*	X*		X*					
<i>Ortochile nigrocoerulea</i> Latreille, 1809	X*	X*	X*	X*	X*						
<i>Poecilobothrus chrysozygos</i> (Wiedemann, 1817)							X				

Table I – continued

Species	Balıkesir	Bilecik	Bursa	Çanakkale	Edime	İstanbul	Kırklareli	Kocaeli	Sakarya	Tekirdağ	Yalova
<i>P. nobilitatus</i> (Linneaus, 1767)				X*							
<i>P. principalis</i> (Loew, 1861)			X*					X			
<i>P. regalis</i> (Meigen, 1824)	X*	X*	X*	X*	X	X*	X	X*	X*	X	X*
<i>Sybistroma discipes</i> (Germar, 1821)				X	X						
<i>S. sphenopterus</i> (Loew, 1859)			X*	X			X	X*			
<i>S. transcaucasica</i> Stackelberg, 1941				X							
<i>Tachytrectus genualis</i> Loew, 1857									X		
<i>T. notatus</i> (Stannius, 1831)										X	
<i>Hydrophorus balticus</i> (Meigen, 1824)	X*		X*	X*			X	X*			
<i>H. praecox</i> (Lehmann, 1822)			X*								
<i>Gymnopternus celer</i> (Meigen, 1824)					X						
<i>Liancalus virens</i> (Scopoli, 1763)	X*	X*	X*	X*			X	X*			
<i>Orthoceratium lacustre</i> (Scopoli, 1763)					X*						
<i>Scellus notatus</i> (Fabricius, 1781)	X*		X*	X*			X				
<i>Schoenophilus versutus</i> Haliday, 1851										X	
<i>Neurigona erichsoni</i> Zetterstedt, 1843			X	X							
<i>N. nubifera</i> (Loew, 1869)					X						
<i>Chrysotimus molliculus</i> (Fallén, 1823)							X				
<i>Peloropodes acuticornis</i> (Oldenberg, 1916)							X			X	
<i>Rhaphium appendiculatum</i> Zetterstedt, 1849			X*	X*			X				
<i>R. brevicorne</i> Curtis, 1835										X	
<i>R. calliginosum</i> Meigen, 1824		X*								X	
<i>Sciapus bellus</i> (Loew, 1873)								X*			
<i>Campsicnemus curvipes</i> (Fallén, 1823)			X*				X		X*		
<i>C. magius</i> (Loew, 1845)			X*								
<i>Sympycnus pulicarius</i> (Fallén, 1823)			X*	X*			X				
<i>Syntormon denticulatus</i> (Zetterstedt, 1843)			X*	X*							
<i>S. pallipes</i> (Fabricius, 1794)	X*	X*	X*	X*	X		X	X*	X*	X	
<i>Teuchophorus simplex</i> Mik, 1880							X				
<i>Xanthochlorus tenellus</i> (Wiedemann, 1817)				X*			X		X*		
Total	8	7	24	20	4	1	23	7	11	7	1




Figure 1. The number of already-known and newly-added species by provinces.

Acknowledgements

The author would like to thank the Scientific and Technical Council of Turkey (TUBITAK, Research Projects No TBAG-107T678) for its financial support of fieldwork in the framework of the project entitled 'Investigation on Biodiversity of Tipuloidea (Diptera) Fauna of Marmara Region', Muğla Sıtkı Koçman University (Research Project No: 17/207) for additional support and Dr Hasan Koç, Dr Okan Özgül, Yağız Molla and A. Levent Yıldırım (Turkey) who supplied me with trap samples.

References

- Grichanov, I. Ya., Tonguç, A., Civelek, H. S., Vikhrev, N. E., Özgül, O., & Dursun, O. (2007). Review of the Turkish Dolichopodidae (Diptera) with first description of male *Hercostomus phoebus* Parent, 1927, new synonyms and new records. *Caucasian Entomological Bulletin*, 3(2), 261-268.
- Grichanov, I. Ya. (2017). *Alphabetic list of generic and specific names of predatory flies of the epifamily Dolichopodidae (Diptera) Second edition*. Saint Petersburg: All-Russian Research Institute of Plant Protection Public., (Plant Protection News. Supplements), 561 pp.
- Grichanov, I. Ya., & Tonguç A. (2010a). New contribution to the Turkish Dolichopodidae fauna (Diptera). *Acta Zoologica Bulgarica*, 62(3), 355-357.
- Grichanov I. Ya., & Tonguç A. (2010b). New contribution to the Turkish Dolichopodidae (Diptera) fauna and taxonomy. *International Jornal of Dipterological Research*, 21(3), 225-229.
- Grichanov, I. Ya., & Ahmadi, A. (2017). Palearctic species of the genus *Lamprochromus* Mik, 1878 (Diptera: Dolichopodidae). *Far Eastern Entomologist*, 336, 1-12.

- Kechev, M., Naglis, S., Tonguç, A., & Pollet, M. (2020). A review of the family Dolichopodidae (Diptera, Empidoidea) of the Balkan Peninsula, with first records for Bulgaria, Montenegro, North Macedonia, and for the European part of Turkey. *Zootaxa*, 4819(3), 436-472.
- Küçükberber, M., Tonguç, A., & Koç, H. (2017). Dolichopodidae (Diptera) fauna of Spil Mountain with four new records. *Türkiye entomoloji bülteni*, 7(1), 23-29.
- Naglis, S., & Negrobov, O. P. (2017). West Palaearctic species of the *Hercostomus* species-group III (Diptera, Dolichopodidae), with description of a new species from Turkey. *Alpine Entomology*, 1, 33-38.
- Tonguç, A., Grichanov, I. Ya., & Kechev, M. (2009). New Records of the Family Dolichopodidae (Diptera) from Turkey. *Acta Zoologica Bulgarica*, 61(2), 213-216.
- Tonguç, A., Grichanov, I. Ya., Koç, H., Özgül, O., & Barlas, M. (2010). Contributions to the Dolichopodidae (Diptera) Fauna of Turkey. *Journal of Entomological Research Society*, 12(2), 103-107.
- Tonguç, A., Grichanov, I. Ya., & Naglis, S. (2016a). Checklist of the Dolichopodidae (Diptera, Brachycera) of Turkey. *Turkish Journal of Zoology*, 40, 14-26.
- Tonguç, A., Grootaert, P., & Barlas, M. (2016b). A new species of *Hercostomus* Loew, 1857 (Diptera: Dolichopodidae) from Turkey. *Zoology in the Middle East*, 62(4), 363-366.
- Yang, D., Zhu, Y., Wang, M., & Zhang, L. (2006). *World Catalog of Dolichopodidae (Insecta: Diptera)*. China Agricultural University Press, 704 pp.

ДОПРИНОС ПОЗНАВАЊУ ФАУНЕ DOLICHOPODIDAE (DIPTERA) РЕГИОНА МАРМАРА У ТУРСКОЈ СА НОВИМ НАЛАЗИМА

Алпер Тонгуч

Извод

Теренска истраживања спроведена су у провинцијама Баликесир, Билеџик, Бурса, Чанакале, Едирне, Истанбул, Измит, Киркларели, Сакарија, Текирдаг и Јалова у регији Мармара у Турској у периоду од 2009. до 2010. године. Идентификацијом прикупљених примерака утврђено је 17 родова и 31 врста из подфамилија Diaphorinae, Dolichopodinae, Hydrophorinae, Rhaphininae, Sciapodinae, Sympycninae, Xanthochlorinae. Две врсте (*Hercostomus plagiatus* Loew, 1857 и *Poecilobothrus nobilitatus* Linnaeus, 1767) су нови налази за фауну долихоподида Турске.

Received: July 14th, 2020

Accepted: August 17th, 2020